

AGU: Journal of Geophysical Research, Space Physics

Keywords

- TLEs
- thunderstorm
- aerosols

Index Terms

- Atmospheric Processes: Atmospheric electricity
- Atmospheric Composition and Structure: Middle atmosphere: energy deposition
- Atmospheric Processes: Mesoscale meteorology
- Atmospheric Processes: Clouds and aerosols
- Global Change: Atmosphere

Abstract

JOURNAL OF GEOPHYSICAL RESEARCH, VOL. 115, A00E58, 20 PP., 2010
doi:10.1029/2009JA014857

Observations of prolific transient luminous event production above a mesoscale convective system in Argentina during the Sprite2006 Campaign in Brazil

F. T. São Sabbas

Instituto Nacional de Pesquisas Espaciais, São José dos Campos, São Paulo, Brazil

M. J. Taylor

Center for Atmospheric and Space Sciences, Utah State University, Logan, Utah, USA

P.-D. Pautet

Center for Atmospheric and Space Sciences, Utah State University, Logan, Utah, USA

M. Bailey

Center for Atmospheric and Space Sciences, Utah State University, Logan, Utah, USA

S. Cummer

Electrical and Computer Engineering Department, Duke University, Durham, North Carolina, USA

R. R. Azambuja

Instituto Nacional de Pesquisas Espaciais, São José dos Campos, São Paulo, Brazil

J. P. C. Santiago

Instituto Nacional de Pesquisas Espaciais, São José dos Campos, São Paulo, Brazil

J. N. Thomas

Department of Electrical and Computer Engineering, Digipen Institute of Technology, Redmond, Washington, USA

O. Pinto

Instituto Nacional de Pesquisas Espaciais, São José dos Campos, São Paulo, Brazil

N. N. Solorzano

Department of Electrical and Computer Engineering, Digipen Institute of Technology, Redmond, Washington, USA

N. J. Schuch

Southern Regional Space Research Center, CRS, INPE-MCT, Santa Maria, Rio Grande do Sul, Brazil

S. R. Freitas

Instituto Nacional de Pesquisas Espaciais, São José dos Campos, São Paulo, Brazil

N. J. Ferreira

Instituto Nacional de Pesquisas Espaciais, São José dos Campos, São Paulo, Brazil

J. C. Conforte

Instituto Nacional de Pesquisas Espaciais, São José dos Campos, São Paulo, Brazil

On the night of 22–23 February 2006, 444 transient luminous events (TLEs), 86% sprites, were observed above a prolific mesoscale convective system (MCS) over Argentina, as part of the third sprite campaign in Brazil. GOES infrared (IR) cloud top temperatures (T_c) and Tropical Rainfall Measuring Mission (TRMM) radar (PR) and microwave (TMI) data were used to investigate the MCS convective characteristics and their relationship with World Wide Location Network (WWLLN) detected cloud-to-ground (CG) lightning and TLE activity. The MCS had a minimum lifetime of 20 hours, 8.5 as a MCS, a maximum extent of $\sim 430,000 \text{ km}^2$, and gusty winds of $\sim 39\text{--}50 \text{ km/h}$. It had several distinctive characteristics: exceptionally high TLE rate, multicellular structure with 19 distinguishable convective regions, and cloud tops temperatures (T_c) $\sim 10\text{--}20 \text{ }^\circ\text{C}$ higher than regular TLE-producing MCSs over the central USA and South America. Most TLEs occurred above “individual stratiform regions”, where T_c varied from $-45 \text{ }^\circ\text{C}$ to $-53 \text{ }^\circ\text{C}$ from the beginning to the end of the night, surrounding the areas of strong convections, with convective cores at T_c $-59 \text{ }^\circ\text{C}$ to $-74 \text{ }^\circ\text{C}$, which did not extend up to or overshoot the tropopause, estimated at $-75 \text{ }^\circ\text{C}$ ($\sim 17.1 \text{ km}$) as normally observed for TLE-producing MCS in these regions. The moderated convection is contrary to the expectation that large charge production is accompanied by vigorous updrafts within deep convection that give rise to cold cloud overshooting tops, thus prompting a detailed study of this prolific TLE-producing thunderstorm. On the basis of a charge moment change threshold of 350 Ckm

and estimated 5 km charge removal altitude, a lower threshold of $\sim 4,300$ C/h was estimated for the hourly charge transfer rate necessary for the observed sprite production (383 events), which is twice the rate for an average TLE-producing MCS (70 events), also estimated. TMI/TRMM data for the storm at early development showed a low brightness temperature of 84 K, indicative of significant ice content, which is important for cloud electrification processes. We suggest that the unusually high incidence of TLEs in this moderately convective MCS may be related to other local geophysical phenomena such as a large tropospheric aerosol concentration due to smoke from forest fires. Satellite fire count data showed that there were ~ 200 fires between 20 and 22 February immediately north of the MCS initiation region and a transport simulation with the Coupled Aerosol-Tracer Transport model from the Brazilian developments on Regional Atmospheric Modeling System (CATT/BRAMS) model showed a large $PM_{2.5}$ aerosol concentration, $10,000 \mu\text{g}/\text{m}^2$ (column integrated), at the region where the MCS developed. The aerosols present in the smoke may have been a source of ice nuclei affecting the production of ice particles that get positively charged, accounting for the charge transfer rate necessary to originate the observed TLE production.

Received 1 September 2009; accepted 28 May 2010; published 2 November 2010.

Citation: São Sabbas, F. T., et al. (2010), Observations of prolific transient luminous event production above a mesoscale convective system in Argentina during the Sprite2006 Campaign in Brazil, *J. Geophys. Res.*, 115, A00E58, doi:10.1029/2009JA014857.

Cited By

Please wait one moment ...

©2010. American Geophysical Union. All Rights Reserved.