39th COSPAR Scientific Assembly 2012

Space Studies of the Upper Atmospheres of the Earth and Planets including Reference Atmospheres (C)

Recent Advances in Equatorial, Low- and Mid-Latitude Mesosphere, Thermosphere and Ionosphere Studies (C1.1)

CORRELATION BETWEEN THE COSMIC NOISE ABSORPTION CALCU-LATED FROM SARINET DATA AND THE ENERGETIC PARTICLES MEA-SURED BY MEPED: SIMULTANEOUS OBSERVATIONS OVER SAMA RE-GION

Juliano Moro, juliano@dae.inpe.br

INPE, Sao Jose Dos Campos, Brazil

Clezio Marcos Denardini, denardin@dae.inpe.br

INPE, Sao Jose Dos Campos, Brazil

Emília Correia, ecorreia@craam.mackenzie.br

Instituto Nacional de Pesquisas Espaciais - INPE, São Paulo, Brazil

Mangalathayil A Abdu, maabdu@dae.inpe.br

Instituto Nacional de Pesquisas Espaciais - INPE, Sao Jose Dos Campos, Brazil

Nelson Jorge Schuch, njschuch@lacesm.ufsm.br

Southern Regional Space Research Center - CRS/CCR/INPE - MCTI, Santa Maria - RS, Brazil

Kazuo Makita, kmakita@la.takushoku-u.ac.jp

Takushoku University, Tokyo, Japan

Laysa Cristina Araujo Resende, laysa@dae.inpe.br

INPE, Sao Jose Dos Campos, Brazil

Lais Maria Guizelli, lais@dae.inpe.br

INPE, Sao Jose Dos Campos, Brazil

Sony Su Chen, sony.chen@dae.inpe.br

INPE, Sao Jose Dos Campos, Brazil

The cosmic noise absorption (CNA) studies are presented in terms of the CNA images obtained from the imaging riometers operated at the Southern Space Observatory, in Sao Martinho da Serra (29.4° S, 53.1° W), Brazil, Concepcion (36.5° S, 73.0° W) and Punta Arenas (53.0° S, 70.5° W) in Chile, which form part of the South American Riometer Network (SARINET) and are located at the central and periphery regions of the South American Magnetic Anomaly (SAMA). Correlations are made between the maximum CNA observed at the SARINET stations and the two channels of energetic electrons (>30 keV and >300 keV) and the three channels of energetic protons (80-240 keV, 800-2500 keV and >6900 keV) from Medium Energy Proton and Electron Detector (MEPED), measured during the moderate geomagnetic storm that occurred on September 3, 2008. The results show high correlations between the CNA detected at SSO and

the flux of protons with energy between 80 and $240~\rm keV$, and the flux of electrons with energies higher than $300~\rm keV$, while an additional ionization at CON was correlated with electrons of energies higher than $30~\rm keV$. The CNA detected at PAC was probably caused by the increase of the flux of protons with energy between $80~\rm and~240~\rm keV$.