

IN13A-23: OPTIMIZATION FIREFLY METHOD FOR WEIGHTED ENSEMBLE OF CONVECTIVE PARAMETERIZATIONS. PART II: SENSITIVITY EXPERIMENT USING TRMM SATELLITE DATA

Ariane Frassoni dos Santos¹, Saulo R. Freitas¹, Eduardo F. P. Luz²,

Haroldo F. de Campos Velho², Manoel A. Gan¹

¹Center for Weather Prediction and Climate Studies, National Institute for Space Research, Cachoeira Paulista, SP

²Laboratory of Computing and Applied Mathematics, National Institute for Space Research,

São José dos Campos, SP

ariane.frassoni@cptec.inpe.br

ABSTRACT

The inverse problem methodology for parameter estimation is applied to a meteorological phenomenon that causes intense rainfall over South America. It is formulated as an optimization problem, where the goal is to apply the Firefly method (FA) as an optimizer for retrieving the weights of the ensemble of convective parameterizations of Grell and Dévényi. The forward problem is the precipitation field of the ensemble of convective parameterizations expressed by several methodologies. The precipitation fields were used as the direct problem (see companion paper), and the precipitation field estimated by the Tropical Rainfall Measuring Mission (TRMM) satellite as the observed data. The inverse problem is solved as an optimization problem with regularization operator of Tikhonov of zero order.

INTRODUCTION

Parameterization of convection

Many different approaches exist concern to:

- What do you use to decide where convection will form
- How strong it will be (*closure*)
- What is important with respect to how convection modulates the environment (*feedback*)

Ensemble version of convective parameterization

- Build more stochasticism into parameterization
- Allows to find objective ways to determine weighting of ensembles for feedback

FIREFLY METHOD

Pseudo code

```

begin
  Objective function f(x),   x=(x1, ..., xd)T
  Generate initial population of fireflies xi (i=1, 2, ..., n)
  Light intensity Ii at xi is determined by f(xi)
  Define light absorption coefficient γ
  while (t < MaxGeneration) (Number of iterations)
    for i = 1 : n all n fireflies
      for j = 1 : d loop over all d dimensions
        if (Ij > Ii), Move firefly i towards j: end if
        Attractiveness varies with distance r via exp[-γr]
        evaluate new solutions and update light intensity
      end for j
    end for i
    Rank the fireflies and find the current best
  end while
  Postprocess results and visualization
end
```

Adapted of Yang (2008)

Light intensity **In a simplest form**

$$I \propto f \propto \frac{I_{\text{fonte}}}{r^2}$$

Movement of the firefly i toward firefly j (brightest)

$$x_i = x_i + \frac{\beta_0}{1 + r^2 \gamma} (x_j - x_i) + \alpha [rand - \frac{1}{2}]$$



attraction randomness

$\gamma = O(1) \Rightarrow$ determines the convergence velocity

To an environment light absorption coefficient fix γ

$$I = I_0 e^{-\gamma r} \quad \rightarrow \quad I = I_0 e^{-r^2 \gamma} \quad \rightarrow \quad I_r = \frac{I_{\text{fonte}}}{1 + r^2 \gamma}$$

RESULTS



AKNOWLEDGEMENTS


The first author aknowlege the research support foundation in the state of São Paulo (Fundação de Amparo à Pesquisa do Estado de São Paulo, FAPESP)

REFERENCES

- Grell, G. A., and Dévényi, D. A generalized approach to parameterizing convection combining ensemble and data assimilation techniques. Geophys. Res. Lett., v. 29, no. 14, 2002
Luz, E. F. P.; Becceneri, J. C.; de Campos Velho, H. F. Conceitualização do algoritmo vagalume e sua aplicação na estimativa de condição inicial de calor. In: IX Workshop do Curso de Computação Aplicada do INPE, 2009
Yang, X. Nature-Inspired Metaheuristic Algorithms, Cambridge, 2008

METHODOLOGY

The model CATT-BRAMS



Cumulus Parametrization of de Grell & Dévényi Multidimensional ensemble

- ✓ Trigger function (dimension 3)
- ✓ Precipitation efficiency (dimension 3)
- ✓ Closures (dimension 5 x 3):
 - Grell, 1993
 - Arakawa & Schubert, 1974
 - Kain e Fritsch, 1993
 - Moisture convergence (tipo Kuo 1965, 1974)
 - Low-level omega (Brown 1979, Frank e Cohen 1987)

Weights estimation – Inverse Problems

Real experimental data (TRMM)

$$J[P] = \sum_{i=1}^W [P_M - P_{TRMM}]^2 \quad \text{where} \quad P_M = \sum_i w_i P_i$$

Numerical experiment: the use of the firefly algorithm, with different number of iterations and number of fireflies

Iterations (MaxGeneration)= 10
Nº fireflies (n) = 5

B₀= 1
α = 0,2
γ = 1

CONCLUSION

The retrieved field of precipitation was in agreement with the observed field. We computed the error precipitation field obtained with simple ensemble average and the error to the retrieved precipitation: errors of the ensemble average are greater than the errors of the retrieved precipitation. We expect to employ the method introduced here to improve the simulated precipitation of the CCATT-BRAMS system.